

Future Computational Needs for Climate Change Modeling

Michael Wehner

Lawrence Berkeley National Laboratory

mfwehner@lbl.gov

510-495-2527



NATIONAL ENERGY RESEARCH
SCIENTIFIC COMPUTING CENTER

Advancing Computational Science of Scale—
Producing Real Results



Background

- ✍ Modeling climate change is high profile science
 - ✍ Politically and socially relevant
 - ✍ Often covered in the popular press
- ✍ Scientific progress is limited by observational and computational constraints.
- ✍ The complexity of the natural climate system must be reflected in numerical models if we hope to gain understanding.

General Circulation Models (GCMs)- Present

- ✍ Fully coupled GCMs are our most sophisticated computational tools.
- ✍ The current state of the art includes submodels of:
 - ✍ Atmosphere
 - ✍ Ocean
 - ✍ Sea ice
 - ✍ Land processes
- ✍ Essentially a fancy hydrodynamics model with thermodynamic source terms.
 - ✍ Tracks energy, momentum and moisture.



NATIONAL ENERGY RESEARCH
SCIENTIFIC COMPUTING CENTER

Advancing Computational Science of Scale—
Producing Real Results



General Circulation Models (GCMs)-Future

- ✍ Current developmental models also target chemical processes.
- ✍ The most important of these from a climate change perspective is biological.
 - ✍ Add carbon and other nutrients to the prognosis
- ✍ Atmospheric chemistry adds numerous other prognostic variables.
- ✍ All of this will further add to the computational burden.



NATIONAL ENERGY RESEARCH
SCIENTIFIC COMPUTING CENTER

Advancing Computational Science of Scale—
Producing Real Results



General Circulation Models (GCMs) Limitations

- ✂ GCMs are expensive.
 - ✂ They exhibit poor scalability in several respects.
- ✂ Example: Atmospheric dynamics
 - ✂ Stable finite difference solution of the Navier-Stokes equations is limited by the Courant Condition.
 - ✂ $\Delta t < \Delta x / v$
 - ✂ Δx = grid spacing, v = maximum wind speed
 - ✂ The consequence of this condition is that computational burden increases nonlinearly as resolution is increased.



NATIONAL ENERGY RESEARCH
SCIENTIFIC COMPUTING CENTER

Advancing Computational Science of Scale—
Producing Real Results



General Circulation Models (GCMs) Limitations

- ✍ Climate change integrations must integrate for multiple centuries.
- ✍ Courant conditions are measured in minutes.
- ✍ This large number of time steps is the principal limiting factor in climate modeling.
 - ✍ Prevents the exploitation of large numbers of processors.
- ✍ Example.
 - ✍ Domain decomposition of grid based hydrodynamics schemes requires that the ratio of “interior” cells to “border” cells be high for parallel efficiency
 - ✍ Doubling the horizontal resolution = Four times as many cells
 - ✍ Four times as many processors can be used at the same efficiency
 - ✍ But each processor has twice as much work to do. Time step was halved. Run time is twice as long.







NATIONAL ENERGY RESEARCH
SCIENTIFIC COMPUTING CENTER

Advancing Computational Science of Scale—
Producing Real Results




Community Climate System Model (NCAR)

Current resolution

-  Atmosphere: Grid=300km. $\Delta t=20$ minutes
-  Ocean: Grid=60km $\Delta t=60$ minutes
-  one simulated year takes five machine hours (128PE SP3)
-  1000 year control run integration required 9 calendar months (7 charged months)

Sounds bad, but this is really good!

-  1990 AMIP1: Many modeling groups required a calendar year to complete a 10 year integration of a stand alone atmospheric general circulation model. Typical resolution was 600km or more.





NATIONAL ENERGY RESEARCH
SCIENTIFIC COMPUTING CENTER

Advancing Computational Science of Scale—
Producing Real Results






What we really want.

Atmosphere

-  Regional climate change prediction will require horizontal grid resolution of 10km (3600X1800)
-  Cloud physics parameterizations could exploit 100 vertical layers

Ocean

-  Mesoscale (~50km) eddies are thought to be crucial to ocean heat transport
-  0.1° grid will resolve these eddies (3600X1800)
-  Short stand-alone integrations are underway now.

Ensembles of integrations are required to address issues of internal (chaotic) variability.

-  Current practice is to make 4 realizations. 10 is better.

Time steps will be measured in seconds.



NATIONAL ENERGY RESEARCH
SCIENTIFIC COMPUTING CENTER


Advancing Computational Science of Scale—
Producing Real Results



Simulated precipitation as a function of resolution

Duffy, et al

DJF Precipitation



NATIONAL ENERGY RESEARCH
SCIENTIFIC COMPUTING CENTER

Advancing Computational Science of Scale—
Producing Real Results



What kind of machine do we need?

✂ What we have is:

- ✂ seaborg.nersc.gov: 6080 processor IBM Power 3
- ✂ cheetah.ornl.gov: 864 processor IBM Power4
- ✂ bluesky.ucar.edu 1024 processor IBM Power4
- ✂ blackforest.ucar.edu: 1172 processor IBM Power 3

✂ Typical CCSM2 configuration uses 128 processors

- ✂ 32 atmosphere (OpenMP and MPI)
- ✂ 48 ocean (MPI only)
- ✂ 8 land (OpenMP and MPI)
- ✂ 32 ice (MPI only)
- ✂ 8 coupler (OpenMP only)

✂ No significant speedup for additional processors.

✂ Achieves about 5% of peak on Power 3



NATIONAL ENERGY RESEARCH
SCIENTIFIC COMPUTING CENTER

Advancing Computational Science of Scale—
Producing Real Results



What kind of machine do we need?

- ✍ Climate codes are highly vectorizable.
 - ✍ High resolution atmospheric portion achieves 22% of peak on 8 processors of NEC SX6 (14GFlops)
 - ✍ Turnaround is 3 to 6 times faster than the best I can do on Power3.



NATIONAL ENERGY RESEARCH
SCIENTIFIC COMPUTING CENTER

Advancing Computational Science of Scale—
Producing Real Results



What kind of machine do we need?

- ✍ Two types of runs:
 - ✍ A single 1000 year control run to estimate noise.
 - ✍ Ensemble of 4-10 transient runs to estimate signal.
 - ✍ I.e. simulation of historical record 1870-2002 or a prediction of the future climate.
 - ✍ This is an additional embarrassingly parallel dimension.



NATIONAL ENERGY RESEARCH
SCIENTIFIC COMPUTING CENTER

Advancing Computational Science of Scale—
Producing Real Results



What kind of machine do we need?

- ✗ Seaborg is the wrong kind of machine.
 - ✗ queue is tuned to large node jobs.
 - ✗ Justifiably so due to large investment in the network.
 - ✗ ensemble runs are made individually. Jobs are still small despite overall usage of more nodes.
 - ✗ 5% of peak?
 - ✗ Too slow to exploit large numbers of processors.



NATIONAL ENERGY RESEARCH
SCIENTIFIC COMPUTING CENTER

Advancing Computational Science of Scale—
Producing Real Results



What kind of machine do we need?

- ✍ A machine with faster processors allows the usage of more processors for a single realization. Remember the courant condition.
 - ✍ Allows higher resolution with more processors at comparable parallel efficiency (grid cells per PE)
- ✍ Example: Increase atmospheric resolution from 300km to 10km.
 - ✍ If you increase the number of processors by 30^2 (900) to maintain the same parallel efficiency (~200 cells per processor) and
 - ✍ If you increase sustained per processor performance by a factor of 30 to compensate for the time step reduction then
 - ✍ ***Turnaround time stays the same.***
 - ✍ In other words, ~30000 processors each achieving ~2GFlops sustained. Six Earth Simulators.



NATIONAL ENERGY RESEARCH
SCIENTIFIC COMPUTING CENTER

Advancing Computational Science of Scale—
Producing Real Results



Conclusions

- ✍ Despite large strides in computing ability, climate change prediction needs are not yet being met.
- ✍ Limitations are imposed by per processor sustained speed.
- ✍ Machines large numbers of processors could be exploited better but only when processors get much faster.
 - ✍ Large numbers of processors are not bad. Slow processors are bad.